Homework No. 01 (Spring 2018)

PHYS 510: Classical Mechanics

Date: 2018 Jan 30

1. (**20 points.**) Given

$$F[u] = \int_{x_1}^{x_2} dx \, a(x) \frac{du(x)}{dx}.$$
 (1)

Evaluate

(a)

(b)

(c)

$$\frac{\delta F}{\delta u(x)}.$$
(2)

2. (20 points.) (Gelfand and Fomin, Calculus of Variations.) Evaluate the functional derivatives of the following functionals, assuming no variation at the end points.

$$F[y] = \int_0^1 dx \, \frac{dy}{dx} \tag{3}$$

$$F[y] = \int_0^1 dx \, y \frac{dy}{dx} \tag{4}$$

$$F[y] = \int_0^1 dx \, xy \frac{dy}{dx} \tag{5}$$

(d)

$$F[y] = \int_{a}^{b} \frac{dx}{x^{3}} \left(\frac{dy}{dx}\right)^{2} \tag{6}$$

3. (20 points.) The electrostatic energy of a charge distribution $\rho(\mathbf{r})$ is

$$E[\rho] = \frac{1}{2} \int d^3r \int d^3r' \frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}.$$
(7)

Evaluate

$$\frac{\delta^2 E}{\delta \rho(\mathbf{r}) \rho(\mathbf{r}')}.$$
(8)

4. (20 points.) The distance between two points in a plane is characterized by the infinitesimal statement

$$ds^2 = dx^2 + dy^2. (9)$$

Let us prove the intuitively obvious statement that the curve of shortest distance going through two points (x_1, y_1) and (x_2, y_2) in a plane, the geodesics of a plane, is a straight line passing through the two points. This amounts to finding the extremal of the functional

$$l[y] = \int_{(x_1,y_1)}^{(x_2,y_2)} ds = \int_{(x_1,y_1)}^{(x_2,y_2)} dx \sqrt{1 + \left(\frac{dy}{dx}\right)^2}.$$
 (10)

Using the fact that we require the curve to necessarily pass through the points (x_1, y_1) and (x_2, y_2) , show that

$$\frac{\delta l[y]}{\delta y(x)} = -\frac{d}{dx} \left[\frac{\frac{dy}{dx}}{\sqrt{1 + \left(\frac{dy}{dx}\right)^2}} \right].$$
(11)

Then, using the condition that the functional derivative is zero for the extremal curve, derive the equation of the geodesic to be given by

$$\frac{dy}{dx} = c,\tag{12}$$

where c is a contant. Identify this as the equation of a straight line in a plane. Find c.