

Homework No. 08 (Fall 2018)

PHYS 520A: Electromagnetic Theory I

Due date: Friday, 2018 Dec 7, 4.00pm

1. **(50 points.)** Let the space be filled with two dielectric materials, with a discontinuity at $z = 0$, such that

$$\varepsilon(\mathbf{r}) = \varepsilon_2 \theta(-z) + \varepsilon_1 \theta(z), \quad (1)$$

where

$$\varepsilon_2 < \varepsilon_1. \quad (2)$$

In addition there is a point charge at q at

$$\mathbf{r}' = 0 \hat{\mathbf{x}} + 0 \hat{\mathbf{y}} + a \hat{\mathbf{z}}. \quad (3)$$

In the following we shall determine the electric potential and electric field everywhere for this configuration.

- (a) Starting from the Maxwell equations (in vacuum) the electric potential for a single point charge at \mathbf{r}' is

$$-\varepsilon_0 \nabla^2 \phi(\mathbf{r}) = q \delta^{(3)}(\mathbf{r} - \mathbf{r}'). \quad (4)$$

Construct the corresponding Green's function to satisfy

$$-\varepsilon_0 \nabla^2 G_0(\mathbf{r} - \mathbf{r}') = \delta^{(3)}(\mathbf{r} - \mathbf{r}'), \quad (5)$$

which is obtained by replacing the charge density with that of a point charge of unit magnitude that is achieved by simply choosing $q = 1$ in this case. The electric potential is in general given in terms of the Green's function by the superposition principle

$$\phi(\mathbf{r}) = \int d^3 r' G_0(\mathbf{r}, \mathbf{r}') \rho(\mathbf{r}'), \quad (6)$$

which for a point charge in vacuum simply reads

$$\phi(\mathbf{r}) = q G_0(\mathbf{r}, \mathbf{r}'), \quad G_0(\mathbf{r}, \mathbf{r}') = \frac{1}{4\pi\varepsilon_0} \frac{1}{|\mathbf{r} - \mathbf{r}'|}, \quad (7)$$

Observe that the configuration under consideration, a point charge in a planar dielectric region, has translation symmetry in x and y directions. Thus, we use Fourier transformation in these coordinates to write

$$G_0(\mathbf{r}, \mathbf{r}') = \int \frac{d^2 k_\perp}{(2\pi)^2} e^{i\mathbf{k}_\perp \cdot (\mathbf{r} - \mathbf{r}')_\perp} g_0(z, z'; k_\perp), \quad (8)$$

where $g_0(z, z'; k_\perp)$ is the Fourier transform of $G_0(\mathbf{r}, \mathbf{r}')$ in the x and y coordinates. Here the subscript \perp is the projection in the plane perpendicular to $\hat{\mathbf{z}}$. Show that the reduced Green's function $g_0(z, z'; k_\perp)$ satisfies the differential equation

$$-\left(\frac{d^2}{dz^2} - k_\perp^2\right) \varepsilon_0 g_0(z, z'; k_\perp) = \delta(z - z'). \quad (9)$$

Show that

$$g_0(z, z'; k_\perp) = \frac{1}{\varepsilon_0} \frac{1}{2k_\perp} e^{-k_\perp |z - z'|}. \quad (10)$$

Thus, find the identity

$$\begin{aligned} \frac{1}{4\pi} \frac{1}{|\mathbf{r} - \mathbf{r}'|} &= \frac{1}{4\pi} \frac{1}{\sqrt{(x - x')^2 + (y - y')^2 + (z - z')^2}} \\ &= \int \frac{d^2 k_\perp}{(2\pi)^2} e^{i\mathbf{k}_\perp \cdot (\mathbf{r} - \mathbf{r}')_\perp} \frac{1}{2k_\perp} e^{-k_\perp |z - z'|}. \end{aligned} \quad (11)$$

- (b) Starting from the macroscopic Maxwell equations the electric potential for a single point charge at \mathbf{r}' in the presence of a dielectric material is

$$-\nabla \cdot [\varepsilon(\mathbf{r}) \nabla] \phi(\mathbf{r}) = q \delta^{(3)}(\mathbf{r} - \mathbf{r}'). \quad (12)$$

Construct the corresponding Green's function to satisfy

$$-\nabla \cdot [\varepsilon(\mathbf{r}) \nabla] G(\mathbf{r}, \mathbf{r}') = \delta^{(3)}(\mathbf{r} - \mathbf{r}'). \quad (13)$$

Show that the corresponding reduced Green's function $g(z, z'; k_\perp)$ satisfies the differential equation

$$\left[-\frac{\partial}{\partial z} \varepsilon(z) \frac{\partial}{\partial z} + \varepsilon(z) k_\perp^2 \right] g(z, z'; k_\perp) = \delta(z - z'), \quad (14)$$

where

$$\varepsilon(z) = \begin{cases} \varepsilon_2, & z < 0, \\ \varepsilon_1, & 0 < z. \end{cases} \quad (15)$$

Show that

$$g(z, z') = \begin{cases} \frac{1}{\varepsilon_2} \frac{1}{2k_\perp} e^{-k_\perp |z - z'|} + \frac{1}{\varepsilon_2} \frac{1}{2k_\perp} \left(\frac{\varepsilon_2 - \varepsilon_1}{\varepsilon_2 + \varepsilon_1} \right) e^{-k_\perp |z|} e^{-k_\perp |z'|}, & z' < 0, \\ \frac{1}{\varepsilon_1} \frac{1}{2k_\perp} e^{-k_\perp |z - z'|} + \frac{1}{\varepsilon_1} \frac{1}{2k_\perp} \left(\frac{\varepsilon_1 - \varepsilon_2}{\varepsilon_1 + \varepsilon_2} \right) e^{-k_\perp |z|} e^{-k_\perp |z'|}, & 0 < z'. \end{cases} \quad (16)$$

(c) Use the identity in Eq. (11) to show that

$$\phi(\mathbf{r}) = q G(\mathbf{r}, \mathbf{r}') = \frac{1}{4\pi\epsilon_1} \frac{q}{|\mathbf{r} - \mathbf{r}'|} + \frac{1}{4\pi\epsilon_1} \frac{q_{\text{im}}}{|\mathbf{r} - \mathbf{r}'_{\text{im}}|}, \quad (17)$$

where

$$q_{\text{im}} = -q \left(\frac{\epsilon_2 - \epsilon_1}{\epsilon_2 + \epsilon_1} \right) \quad (18)$$

and

$$\mathbf{r}'_{\text{im}} = \begin{cases} \mathbf{r}', & z > 0, \\ \mathbf{r}' - 2a \hat{\mathbf{z}}, & z < 0. \end{cases} \quad (19)$$

Thus, prescribe an algorithm to determine the electric potential for planar dielectrics—a method of images for planar dielectrics.

(d) Determine the electric field to be

$$\mathbf{E}(\mathbf{r}) = -\nabla\phi(\mathbf{r}) = \frac{q}{4\pi\epsilon_1} \frac{\mathbf{r} - \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|^3} + \frac{q_{\text{im}}}{4\pi\epsilon_1} \frac{\mathbf{r} - \mathbf{r}'_{\text{im}}}{|\mathbf{r} - \mathbf{r}'_{\text{im}}|^3}. \quad (20)$$

Draw the electric field lines for $\epsilon_2 < \epsilon_1$, and compare it with the electric field lines for $\epsilon_2 > \epsilon_1$.

(e) By evaluating the ratios

$$\frac{E_x(x, y, +\delta)}{E_x(x, y, -\delta)}, \quad \frac{E_y(x, y, +\delta)}{E_y(x, y, -\delta)}, \quad \frac{E_z(x, y, +\delta)}{E_z(x, y, -\delta)}, \quad (21)$$

determine the boundary conditions satisfied by the electric field lines. This is the Snell's law for the electric field lines. Note that the Snell's law for refraction is expressed in terms of the propagation vector of a plane wave, which is perpendicular to the electric field lines.

(f) A perfect conductor (in the static limit) is a dielectric material with a very high dielectric constant ($\epsilon \rightarrow \infty$). Consider the extreme limit

$$\epsilon_1 < \epsilon_2 \rightarrow \infty \quad (22)$$

and

$$\epsilon_2 < \epsilon_1 \rightarrow \infty. \quad (23)$$

Examine these cases critically. Compare your results with the method of images for perfect conductors.