

# Homework No. 11 (2019 Fall)

## PHYS 301: Theoretical Methods in Physics

Due date: Wednesday, 2019 Dec 4, 10:00 AM, in class

0. Keywords: Legendre polynomials, orthogonality relations, completeness relation.
0. Problems 1, 2, and 8 are to be submitted for assessment. Rest are for practice.

1. **(20 points.)** Using Mathematica (or another graphing tool) plot the Legendre polynomials  $P_l(x)$  for  $l = 0, 1, 2, 3, 4$  on the same plot. Note that  $-1 \leq x \leq 1$ . Based on the pattern you see what can you conclude about the number of roots for  $P_l(x)$ . In Mathematica these plots are generated using the following commands:

```
Plot[{LegendreP[0,x], LegendreP[1,x], LegendreP[2,x], LegendreP[3,x],  
LegendreP[4,x]}, {x, -1, 1}]
```

Compare your plots with those in Wikipedia article on ‘Legendre Polynomials’. While there read the Wikipedia article on Adrien-Marie Legendre and the associated ‘Portrait Debacle’.

2. **(20 points.)** Legendre polynomials are conveniently generated using the relation

$$P_l(x) = \left(\frac{d}{dx}\right)^l \frac{(x^2 - 1)^l}{2^l l!}, \quad (1)$$

where  $-1 \leq x \leq 1$ . Evaluate Legendre polynomials of degree  $l = 0, 1, 2, 3, 4$  in this manner.

3. **(20 points.)** Legendre polynomials  $P_l(x)$  satisfy the relation

$$\int_{-1}^1 dx P_l(x) = 0 \quad \text{for } l \geq 1. \quad (2)$$

Verify this explicitly for  $l = 0, 1, 2, 3, 4$ .

4. **(20 points.)** Legendre polynomials satisfy the differential equation

$$\left[ \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \sin \theta \frac{\partial}{\partial \theta} + l(l+1) \right] P_l(\cos \theta) = 0. \quad (3)$$

Verify this explicitly for  $l = 0, 1, 2, 3, 4$ .

5. **(20 points.)** Legendre polynomials satisfy the orthogonality relation

$$\int_{-1}^1 dx P_l(x) P_{l'}(x) = \frac{2}{2l+1} \delta_{ll'}. \quad (4)$$

Verify this explicitly for  $l = 0, 1, 2$  and  $l' = 0, 1, 2$ .

6. **(20 points.)** Legendre polynomials satisfy the completeness relation

$$\sum_{l=0}^{\infty} \frac{2l+1}{2} P_l(x) P_l(x') = \delta(x - x'). \quad (5)$$

This is for your information. No work needed.

7. **(Example.)** The Legendre polynomials of order  $l$  are

$$P_l(x) = \left( \frac{d}{dx} \right)^l \frac{(x^2 - 1)^l}{2^l l!}. \quad (6)$$

In particular,

$$P_0(x) = 1, \quad (7a)$$

$$P_1(x) = x, \quad (7b)$$

$$P_2(x) = \frac{3}{2}x^2 - \frac{1}{2}. \quad (7c)$$

The expansion

$$F(x, t) = \frac{1}{\sqrt{1 - 2xt + t^2}} = \sum_{l=0}^{\infty} t^l P_l(x), \quad |t| < 1, \quad (8)$$

is usually referred to as the generating function for Legendre's polynomials. From it all the properties of these polynomials may be derived.

8. **(Example.)** The Legendre polynomials of order  $l$  satisfy the recurrence relation

$$(2l+1)xP_l(x) = (l+1)P_{l+1}(x) + lP_{l-1}(x), \quad l = 1, 2, 3, \dots \quad (9)$$

Recall,

$$P_0(x) = 1, \quad (10a)$$

$$P_1(x) = x. \quad (10b)$$

Derive the explicit expression for  $P_l(x)$  for  $l = 2, 3, 4, 5$  using the recurrence relation.