

(Preview of) Midterm Exam No. 02 (Fall 2019)

PHYS 500A: Mathematical Methods

Date: 2019 Oct 29

1. (20 points.) On contour integrals with poles.
2. (20 points.) Consider the integral

$$I(\theta) = \frac{1}{\pi} \int_0^\infty \frac{x^{\frac{1}{2}} dx}{1 + 2x \cos \theta + x^2}, \quad (1)$$

where $0 \leq \theta < 2\pi$ is real. To evaluate $I(\theta)$ let us consider the following integral on the complex plane

$$G(\theta) = \frac{1}{\pi} \oint_c \frac{z^{\frac{1}{2}} dz}{1 + 2z \cos \theta + z^2}, \quad (2)$$

where the contour c is described in Figure 1.




Figure 1: Contour $c = c_1 + c_2 + c_3 + c_4$. The radii of the contours c_2 and c_4 are R and ϵ , respectively, and contours c_1 and c_3 are δ away from the real line. We assume limits $\epsilon \rightarrow 0$, $R \rightarrow \infty$, and $\delta \rightarrow 0$.

(a) Show that

$$1 + 2z \cos \theta + z^2 = (z + e^{i\theta})(z + e^{-i\theta}) \quad (3)$$

and identify the poles. Show that the integrand has a branch point at $z = 0$. Choose the branch cut to be the positive real line. Using Cauchy's theorem show that

$$G(\theta) = 2 \frac{\sin \frac{\theta}{2}}{\sin \theta}. \quad (4)$$

(b) Next, let us evaluate $G(\theta)$ by evaluating the integrals on the contour explicitly.

- i. For the part of contour constituting c_1 substitute $z = xe^{i\delta} \sim x + i\delta'$ and show that

$$\lim_{\delta \rightarrow 0} \lim_{\epsilon \rightarrow 0} \lim_{R \rightarrow \infty} \frac{1}{\pi} \oint_{c_1} \frac{z^{\frac{1}{2}} dz}{(z + e^{i\theta})(z + e^{-i\theta})} = I(\theta). \quad (5)$$

- ii. For the part of contour constituting c_3 substitute $z = xe^{i(2\pi-\delta)} \sim x - i\delta'$ and show that

$$\lim_{\delta \rightarrow 0} \lim_{\epsilon \rightarrow 0} \lim_{R \rightarrow \infty} \frac{1}{\pi} \oint_{c_3} \frac{z^{\frac{1}{2}} dz}{(z + e^{i\theta})(z + e^{-i\theta})} = -e^{i\frac{2\pi}{2}} I(\theta) = I(\theta). \quad (6)$$

- iii. For the part of contour constituting c_2 substitute $z = Re^{i\theta}$ and show that

$$\lim_{R \rightarrow \infty} \frac{1}{\pi} \oint_{c_2} \frac{z^{\frac{1}{2}} dz}{(z + e^{i\theta})(z + e^{-i\theta})} = 0. \quad (7)$$

- iv. For the part of contour constituting c_4 substitute $z = \epsilon e^{i\theta}$ and show that

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\pi} \oint_{c_4} \frac{z^{\frac{1}{2}} dz}{(z + e^{i\theta})(z + e^{-i\theta})} = 0. \quad (8)$$

(c) Together, conclude that

$$2 \frac{\sin \frac{\theta}{2}}{\sin \theta} = I(\theta) + 0 + I(\theta) + 0. \quad (9)$$

Thus, evaluate $I(\theta)$.

3. **(20 points.)** On damped harmonic oscillator.
4. **(20 points.)** Read the article titled 'Life at low Reynolds number' by E. M. Purcell, American Journal of Physics 45 (1977) 3. Here is the link to the article:

<http://dx.doi.org/10.1119/1.10903>

One question will be asked to verify the understanding of the concept being discussed in the paper.