

(Preview of) Midterm Exam No. 01 (2020 Spring)

PHYS 510: CLASSICAL MECHANICS

Department of Physics, Southern Illinois University–Carbondale

Date: 2019 Feb 18

1. (20 points.) On functional derivative.
2. (20 points.) Not available in preview form.
3. (20 points.) Not available in preview form.
4. (20 points.) On Lagrangian.
5. (20 points.) Fermat's principle in ray optics states that a ray of light takes the path of least time between two given points. The speed of light in a medium is given in terms of the refractive index

$$n = \frac{c}{v}, \quad (1)$$

of the medium, where c is the speed of light in vacuum and v is the speed of light in the medium. Consider a ray of light traversing a path from (x_1, y_1) to (x_2, y_2) in a plane of fixed z .




Figure 1: Problem 5.

- (a) Show that the time taken to travel an infinitesimal distance ds is given by

$$dt = \frac{ds}{v} = \frac{n ds}{c}, \quad (2)$$

where ds in a plane is characterized by the infinitesimal statement

$$ds^2 = dx^2 + dy^2. \quad (3)$$

- (b) Fermat's principle states that the path traversed by a ray of light from (x_1, y_1) to (x_2, y_2) is the extremal of the functional

$$T[y] = \frac{1}{c} \int_{(x_1, y_1)}^{(x_2, y_2)} n ds = \frac{1}{c} \int_{x_1}^{x_2} dx n(x) \sqrt{1 + \left(\frac{dy}{dx}\right)^2}. \quad (4)$$

- (c) Since the ray of light passes through the points (x_1, y_1) and (x_2, y_2) , we do not consider variations at these (end) points. Thus, show that

$$\frac{\delta T[y]}{\delta y(x)} = -\frac{d}{dx} \left[\frac{n(x) \frac{dy}{dx}}{\sqrt{1 + \left(\frac{dy}{dx}\right)^2}} \right]. \quad (5)$$

- (d) Using Fermat's principle show that the differential equation for the path $y(x)$ traversed by the ray of light is

$$\frac{n(x) \frac{dy}{dx}}{\sqrt{1 + \left(\frac{dy}{dx}\right)^2}} = n_0, \quad (6)$$

where n_0 is a constant. Show that the above equation can be rewritten in the form

$$\frac{dy}{dx} = \frac{n_0}{\sqrt{n(x)^2 - n_0^2}}. \quad (7)$$

- (e) Let us consider a medium with refractive index $(x_1 = a)$

$$n(x) = \begin{cases} \frac{a}{x}, & 0 < x < a, \\ 1, & a < x. \end{cases} \quad (8)$$

Solve the corresponding differential equation to obtain

$$y(x) - y_0 = \frac{1}{n_0} \left[\sqrt{a^2 - n_0^2 x^2} - \sqrt{a^2 - n_0^2 a^2} \right], \quad x < a. \quad (9)$$

The path in this medium satisfies the equation of a circle. Determine the radius of the circle to be a/n_0 and the location of the center to be $(0, y_0 - a\sqrt{(1/n_0^2) - 1})$. For initial conditions

$$y(x_1) = y_1 \quad \text{and} \quad \left. \frac{dy}{dx} \right|_{x=x_1} = y'_1 \quad (10)$$

show that the integration constants are determined to be

$$y_0 = y_1 \quad \text{and} \quad n_0 = \frac{y_1'}{\sqrt{1 + y_1'^2}}. \quad (11)$$

For the special case when $y_1 = 0$ and $y_1' \rightarrow \infty$ show that $n_0 = 1$ and

$$y(x) = \sqrt{a^2 - x^2}, \quad x < a. \quad (12)$$

Evaluate the total time taken for light to go from $(x_1 = a, y_1 = 0)$ to $(x_2 = 0, y_2 = a)$.