

# Homework No. 03 (2020 Spring)

## PHYS 510: CLASSICAL MECHANICS

*Department of Physics, Southern Illinois University–Carbondale*

Due date: Tuesday, 2020 Feb 11, 4.30pm

1. **(60 points.)** Consider a rope of uniform mass density  $\lambda = dm/ds$  hanging from two points,  $(x_1, y_1)$  and  $(x_2, y_2)$ , as shown in Figure 1. The gravitational potential energy of




Figure 1: Problem 1.

an infinitely tiny element of this rope at point  $(x, y)$  is given by

$$dU = dm gy = \lambda g ds y, \quad (1)$$

where

$$ds^2 = dx^2 + dy^2. \quad (2)$$

A catenary is the curve that the rope assumes, that minimizes the total potential energy of the rope.

- (a) Show that the total potential energy  $U$  of the rope hanging between points  $x_1$  and  $x_2$  is given by

$$U[x] = \lambda g \int_{(x_1, y_1)}^{(x_2, y_2)} y ds = \lambda g \int_{y_1}^{y_2} dy y \sqrt{1 + \left(\frac{dx}{dy}\right)^2}. \quad (3)$$

- (b) Since the curve passes through the points  $(x_1, y_1)$  and  $(x_2, y_2)$ , we have no variations at these (end) points. Thus, show that

$$\frac{\delta U[x]}{\delta x(y)} = -\lambda g \frac{d}{dy} \left[ y \frac{\frac{dx}{dy}}{\sqrt{1 + \left(\frac{dx}{dy}\right)^2}} \right]. \quad (4)$$

(c) Using the extremum principle show that the differential equation for the catenary is

$$\frac{dx}{dy} = \frac{a}{\sqrt{y^2 - a^2}}, \quad (5)$$

where  $a$  is an integration constant.

(d) Show that integration of the differential equation yields the equation of the catenary

$$y = a \cosh \frac{x - x_0}{a}, \quad (6)$$

where  $x_0$  is another integration constant.

(e) For the case  $y_1 = y_2$  we have

$$\frac{y_1}{a} = \cosh \frac{x_1 - x_0}{a}, \quad (7a)$$

$$\frac{y_2}{a} = \cosh \frac{x_2 - x_0}{a}, \quad (7b)$$

which leads to the solution, assuming  $x_1 \neq x_2$ ,

$$x_0 = \frac{x_1 + x_2}{2}. \quad (8)$$

Identify  $x_0$  in Figure 1.

(f) Next, derive

$$\frac{y_1}{a} = \frac{y_2}{a} = \cosh \frac{x_2 - x_1}{2a}, \quad (9)$$

which, in principle, determines  $a$ . However, this is a transcendental equation in  $a$  and does not allow exact evaluation of  $a$ , and one depends on numerical solutions. Observe that, if  $x = x_0$  in Eq. (6), then  $y = a$ . Identify  $a$  in Figure 1.

2. (20 points.) A catenary is described by

$$y = a \cosh \left( \frac{x - x_0}{a} \right), \quad (10)$$

where constants  $a$  and  $x_0$  are determined by the position of the end points  $(x_1, y_1)$  and  $(x_2, y_2)$ . Let us choose  $x_0 = 0$  and  $a = 1$  such that

$$y = \cosh x, \quad (11)$$

where  $x$  and  $y$  are dimensionless variables.

(a) Using series expansion show that

$$\cosh x = 1 + \frac{x^2}{2} + \dots \quad (12)$$

(b) The parabola

$$y = 1 + \frac{x^2}{2} \quad (13)$$

is an approximation for the catenary. Plot the above parabola and a catenary in the same plot for  $-1 < x < 1$  and estimate the maximum error in the approximation.