

# Homework No. 11 (2021 Spring)

## PHYS 510: CLASSICAL MECHANICS

*Department of Physics, Southern Illinois University–Carbondale*

Due date: Thursday, 2021 Apr 8, 9.30am

1. (20 points.) (Based on Schwinger, chapter 9) The Hamiltonian for a Kepler problem is

$$H = \frac{p_1^2}{2m_1} + \frac{p_2^2}{2m_2} - \frac{\alpha}{|\mathbf{r}_1 - \mathbf{r}_2|}, \quad (1)$$

where  $\mathbf{r}_1$  and  $\mathbf{r}_2$  are the positions of the two constituent particles of masses  $m_1$  and  $m_2$ .

- (a) Introduce the coordinates representing the center of mass, relative position, total momentum, and relative momentum:

$$\mathbf{R} = \frac{m_1\mathbf{r}_1 + m_2\mathbf{r}_2}{m_1 + m_2}, \quad \mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2, \quad \mathbf{P} = \mathbf{p}_1 + \mathbf{p}_2, \quad \mathbf{p} = \frac{m_2\mathbf{p}_1 - m_1\mathbf{p}_2}{m_1 + m_2}, \quad (2)$$

respectively, to rewrite the Hamiltonian as

$$H = \frac{P^2}{2M} + \frac{p^2}{2\mu} - \frac{\alpha}{r}, \quad (3)$$

where

$$M = m_1 + m_2, \quad \frac{1}{\mu} = \frac{1}{m_1} + \frac{1}{m_2}. \quad (4)$$

- (b) Show that Hamilton's equations of motion are given by

$$\frac{d\mathbf{R}}{dt} = \frac{\mathbf{P}}{M}, \quad \frac{d\mathbf{P}}{dt} = 0, \quad \frac{d\mathbf{r}}{dt} = \frac{\mathbf{p}}{\mu}, \quad \frac{d\mathbf{p}}{dt} = -\frac{\alpha\mathbf{r}}{r^3}. \quad (5)$$

- (c) Verify that the Hamiltonian  $H$ , the angular momentum  $\mathbf{L} = \mathbf{r} \times \mathbf{p}$ , and the Laplace-Runge-Lenz vector

$$\mathbf{A} = \frac{\mathbf{r}}{r} - \frac{\mathbf{p} \times \mathbf{L}}{\mu\alpha}, \quad (6)$$

are the three constants of motion for the Kepler problem. That is, show that

$$\frac{dH}{dt} = 0, \quad \frac{d\mathbf{L}}{dt} = 0, \quad \frac{d\mathbf{A}}{dt} = 0. \quad (7)$$