

Homework No. 05 (2021 Spring)

PHYS 520B: ELECTROMAGNETIC THEORY

Department of Physics, Southern Illinois University–Carbondale

Due date: Tuesday, 2021 Mar 9, 12.30pm

1. (50 points.) Hamiltonian for a charge particle of mass m and charge q in a magnetic field \mathbf{B} is given by

$$H(\mathbf{x}, \mathbf{p}) = \frac{1}{2m} (\mathbf{p} - q\mathbf{A})^2, \quad (1)$$

where

$$\mathbf{B} = \nabla \times \mathbf{A}. \quad (2)$$

Let

$$\frac{\partial \mathbf{A}}{\partial t} = 0. \quad (3)$$

- (a) Show that the Hamilton equations of motion,

$$\frac{d\mathbf{x}}{dt} = \frac{\partial H}{\partial \mathbf{p}}, \quad (4a)$$

$$\frac{d\mathbf{p}}{dt} = -\frac{\partial H}{\partial \mathbf{x}}, \quad (4b)$$

leads to the equations, using ($\mathbf{v} = d\mathbf{x}/dt$)

$$m\mathbf{v} = \mathbf{p} - q\mathbf{A}, \quad (5a)$$

$$\frac{d\mathbf{p}}{dt} = q(\nabla \mathbf{A}) \cdot \mathbf{v}. \quad (5b)$$

Show that the above equations in conjunction imply the familiar equation

$$m \frac{d\mathbf{v}}{dt} = q\mathbf{v} \times \mathbf{B}. \quad (6)$$

- (b) For two functions

$$A = A(\mathbf{x}, \mathbf{p}, t), \quad (7a)$$

$$B = B(\mathbf{x}, \mathbf{p}, t), \quad (7b)$$

the Poisson bracket with respect to the canonical variables \mathbf{x} and \mathbf{p} is defined as

$$[A, B]_{\mathbf{x}, \mathbf{p}}^{\text{P.B.}} \equiv \frac{\partial A}{\partial \mathbf{x}} \cdot \frac{\partial B}{\partial \mathbf{p}} - \frac{\partial A}{\partial \mathbf{p}} \cdot \frac{\partial B}{\partial \mathbf{x}}. \quad (8)$$

Evaluate the Poisson bracket

$$[\mathbf{x}, \mathbf{x}]_{\mathbf{x}, \mathbf{p}}^{\text{P.B.}} = 0. \quad (9)$$

(c) Evaluate the Poisson bracket

$$[\mathbf{x}^i, \mathbf{v}^j]_{\mathbf{x}, \mathbf{p}}^{\text{P.B.}} = \frac{1}{m} \mathbf{1}^{ij}. \quad (10)$$

(d) Evaluate the Poisson bracket

$$[\mathbf{x}^i, \mathbf{p}^j]_{\mathbf{x}, \mathbf{p}}^{\text{P.B.}} = \mathbf{1}^{ij}. \quad (11)$$

(e) Evaluate the Poisson bracket

$$[m\mathbf{v}^i, m\mathbf{v}^j]_{\mathbf{x}, \mathbf{p}}^{\text{P.B.}} = q(\nabla^i \mathbf{A}^j - \nabla^j \mathbf{A}^i). \quad (12)$$

Verify that

$$(\nabla^i \mathbf{A}^j - \nabla^j \mathbf{A}^i) = \varepsilon^{ijk} \mathbf{B}^k = -\mathbf{1} \times \mathbf{B}. \quad (13)$$

Poisson bracket in classical mechanics has direct correspondence to commutation relation in quantum mechanics through the factor $i\hbar$, which conforms with experiments and balances the dimensions. Then, we can write

$$[m\mathbf{v}, m\mathbf{v}] = i\hbar q \mathbf{B} \quad (14)$$

or

$$m\mathbf{v} \times m\mathbf{v} = i\hbar q \mathbf{B}, \quad (15)$$

using the fact that the commutator and the vector product satisfies the same Lie algebra as that of Poisson bracket.