

(Preview of) Midterm Exam No. 02 (Fall 2022)

PHYS 500A: MATHEMATICAL METHODS

School of Physics and Applied Physics, Southern Illinois University–Carbondale

Date: 2022 Nov 4

1. (20 points.) In class.
2. (20 points.) In class.
3. (20 points.) Verify the identity

$$\phi \nabla \cdot (\lambda \nabla \psi) - \psi \nabla \cdot (\lambda \nabla \phi) = \nabla \cdot [\lambda(\phi \nabla \psi - \psi \nabla \phi)], \quad (1)$$

which is a slight generalization of what is known as Green's second identity. Here ϕ , ψ , and λ , are position dependent functions.

4. (20 points.) The expression for the electric potential due to a point charge placed in front of a perfectly conducting semi-infinite slab, described by

$$\frac{\varepsilon(z)}{\varepsilon_0} = \begin{cases} \infty, & z < 0, \\ 1, & 0 < z, \end{cases} \quad (2)$$

is given in terms of the reduced Green function that satisfies the differential equation ($0 < \{z, z'\}$)

$$-\left[\frac{\partial^2}{\partial z^2} - k^2 \right] \varepsilon_0 g(z, z') = \delta(z - z') \quad (3)$$

with boundary conditions requiring the reduced Green's function to vanish at $z = 0$ and at $z \rightarrow \infty$.

- (a) Construct the reduced Green function in the form

$$\varepsilon_0 g(z, z') = \begin{cases} Ae^{kz} + Be^{-kz}, & 0 < z < z', \\ Ce^{kz} + De^{-kz}, & 0 < z' < z, \end{cases} \quad (4)$$

and solve for the four coefficients, A, B, C, D , using the conditions

$$\varepsilon_0 g(0, z') = 0, \quad (5a)$$

$$\varepsilon_0 g(\infty, z') = 0, \quad (5b)$$

$$\varepsilon_0 g(z, z') \Big|_{z=z'-\delta}^{z=z'+\delta} = 0, \quad (5c)$$

$$\partial_z \varepsilon_0 g(z, z') \Big|_{z=z'-\delta}^{z=z'+\delta} = -1. \quad (5d)$$

(b) Express the solution in the form

$$\varepsilon_0 g(z, z') = \frac{1}{2k} e^{-k|z-z'|} - \frac{1}{2k} e^{-k|z|} e^{-k|z'|}. \quad (6)$$

5. (**20 points.**) The expression for the electric potential due to a point charge placed in between two parallel grounded perfectly conducting semi-infinite slabs, described by

$$\frac{\varepsilon(z)}{\varepsilon_0} = \begin{cases} \infty, & z < 0, \\ 1, & 0 < z < a, \\ \infty, & a < z, \end{cases} \quad (7)$$

is given in terms of the reduced Green function that satisfies the differential equation ($0 < \{z, z'\} < a$)

$$\left[-\frac{\partial^2}{\partial z^2} + k^2 \right] \varepsilon_0 g(z, z') = \delta(z - z') \quad (8)$$

with boundary conditions requiring the reduced Green's function to vanish at $z = 0$ and $z = a$.

(a) Construct the reduced Green's function in the form

$$\varepsilon_0 g(z, z') = \begin{cases} A \sinh kz + B \cosh kz, & 0 < z < z' < a, \\ C \sinh kz + D \cosh kz, & 0 < z' < z < a, \end{cases} \quad (9)$$

and solve for the four coefficients, A, B, C, D , using the conditions

$$\varepsilon_0 g(0, z') = 0, \quad (10a)$$

$$\varepsilon_0 g(a, z') = 0, \quad (10b)$$

$$\varepsilon_0 g(z, z') \Big|_{z=z'-\delta}^{z=z'+\delta} = 0, \quad (10c)$$

$$\partial_z \varepsilon_0 g(z, z') \Big|_{z=z'-\delta}^{z=z'+\delta} = -1. \quad (10d)$$

(b) After using conditions in Eqs. (10a) and (10b) show that the reduced Green's function can be expressed in the form

$$\varepsilon_0 g(z, z') = \begin{cases} A \sinh kz, & 0 < z < z' < a, \\ C' \sinh k(a - z), & 0 < z' < z < a, \end{cases} \quad (11)$$

where $C' = -C / \cosh ka$. Then, use Eqs. (10c) and (10d) to show that

$$\varepsilon_0 g(z, z') = \begin{cases} \frac{\sinh kz \sinh k(a - z')}{k \sinh ka}, & 0 < z < z' < a, \\ \frac{\sinh kz' \sinh k(a - z)}{k \sinh ka}, & 0 < z' < z < a. \end{cases} \quad (12)$$

- (c) Take the limit $ka \rightarrow \infty$ in your solution above, (which corresponds to moving the slab at $z = a$ to infinity,) to obtain the reduced Green's function for a single perfectly conducting slab,

$$\lim_{ka \rightarrow \infty} \varepsilon_0 g(z, z') = \frac{1}{2k} e^{-k|z-z'|} - \frac{1}{2k} e^{-k|z|} e^{-k|z'|}. \quad (13)$$

This should serve as a check for your solution to the reduced Green's function. Hint: The hyperbolic functions here are defined as

$$\sinh x = \frac{1}{2}(e^x - e^{-x}) \quad \text{and} \quad \cosh x = \frac{1}{2}(e^x + e^{-x}). \quad (14)$$