

## Homework No. 03 (Fall 2023)

### PHYS 205B: UNIVERSITY PHYSICS

*School of Physics and Applied Physics, Southern Illinois University–Carbondale*

Due date: Tuesday, 2023 Sep 19, 9:30 AM, on D2L

### Instructions

- You are encouraged to use any of the resources to complete this homework. However, the extent to which you depend on resources while doing homework is usually a measure of how much extra work you need to put in to master the associated concepts. Solutions should be the last resource.
- Describe your thought process in detail and organize it clearly. Make sure your answer has units and the right number of significant digits.
- After completion, scan the pages as a single PDF file, and submit the file on D2L (under Assessments → Assignments).

### Problems

1. (10 points.) Consider a region of uniform electric field  $\vec{E} = -E\hat{j}$  of magnitude  $E = 1.0 \times 10^3 \text{ N/C}$  and direction vertically down. Distance between points  $A$  to  $B$  is  $h = 6.0 \text{ cm}$ , and the distance between points  $B$  to  $C$  is  $d = 8.0 \text{ cm}$ . Refer Fig. 1.
  - (a) What is the potential difference between points  $A$  and  $B$ ?
  - (b) What is the potential difference between points  $B$  and  $C$ ?
  - (c) What is the potential difference between points  $A$  and  $C$ ?




Figure 1: Problem 1

### Solution




Figure 2: Problem 2

2. (10 points.) A sphere with uniform charge distribution  $-Q = -3.0 \mu\text{C}$  is fixed at the origin. Point  $A$  is on a sphere of radius  $5.0 \text{ cm}$  and point  $B$  is on a sphere of radius  $10.0 \text{ cm}$ . Refer Figure 2.
- What is the work done by the electric force acting on charge  $q = +2.0 \mu\text{C}$ , when  $q$  is moved from point  $A$  to point  $B$ .
  - What is the change in the electric potential energy between  $-Q$  and  $q$  when  $q$  is moved from point  $A$  to point  $B$ .
  - If there are no other forces acting on charge  $q$ , using the work-energy theorem calculate the change in kinetic energy of charge  $q$ .

### Solution

3. (10 points.) Find the electric potential at the point marked  $\times$  on the  $x$  axis in Figure 3. Given  $q_1 = -1.0 \text{ nC}$ ,  $q_2 = +2.0 \text{ nC}$ ,  $s = 2a$ ,  $t = 3a$ ,  $y = 4a$ ,  $a = 1.8 \text{ cm}$ .




Figure 3: Problem 3

### Solution [2023 Spring, MT-01, Problem 7]

4. (10 points.) Four charges  $q_1 = q$ ,  $q_2 = -2q$ ,  $q_3 = -3q$ , and  $q_4 = 4q$ , are placed at the corners of a square of side  $L$ , such that  $q_1$  and  $q_4$  are at diagonally opposite corners. Refer Figure 4.




Figure 4: Problem 4

- What is the electric potential at the center of square?
- What is the electric potential at point  $a$ ?
- What is the electric potential at point  $b$ ?
- What is the electric potential difference between points  $a$  and  $c$ ?
- How much potential energy is required to move another charge  $q$  from infinity to the center of the square?
- How much additional potential energy is required to move this charge from the center of the square to point  $a$ ?

**Solution**

- (10 points.) Determine the total energy required to assemble four identical positive charges  $Q$  at the corners of a square of length  $L$ . Assume that the charges are brought from infinity.

**Solution**

- (10 points.) A positive charge  $Q_1 = 1.0 \text{ nC}$  is held fixed. Another positive charge  $Q_2 = 2Q_1$  is tied to charge  $Q_1$  using a string of length  $a = 5.0 \text{ cm}$ . Assume the radius of the two charges to be small in comparison to  $a$ . The charges have masses  $m_1 = 0.05 \text{ grams}$  and  $m_2 = 2m_1$ . When the string is cut the two charges fly off in opposite directions. Determine the speed of each of the charges when they are (infinitely) far apart. (Hint: Use conservation of momentum and conservation of energy.)

**Solution**

- (10 points.) Consider a uniformly charged disc of radius  $R$  with charge per unit area  $\sigma$  placed on the  $yz$  plane such that the origin is the center of the disc. Determine the electric potential on the  $x$  axis to be

$$V(x) = -\frac{\sigma}{2\epsilon_0} \left[ x - \sqrt{x^2 + R^2} \right]. \quad (1)$$

Using the fact that the electric field is the negative gradient of the electric potential, calculate the  $x$ -component of the electric field on the  $x$  axis to be

$$E_x = -\frac{\partial V}{\partial x} = \frac{\sigma}{2\epsilon_0} \left[ 1 - \frac{x}{\sqrt{x^2 + R^2}} \right]. \quad (2)$$

- (a) Show that this leads to the potential and electric field of a point charge in the limit  $R \ll x$ .
- (b) Analyze the limit  $x \ll R$ . Show that this leads to the potential and electric field for a non-conducting uniformly charged plate. Plot the electric potential as a function of  $x$  for this case.

**Solution**

8. (10 points.) The electric field inside and outside a conducting sphere of radius  $R$  is given by

$$\mathbf{E} = \begin{cases} 0, & r < R \text{ (inside),} \\ \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \hat{\mathbf{r}}, & R < r \text{ (outside).} \end{cases} \quad (3)$$

Determine the electric potential inside (and outside) the sphere using

$$\Delta V = - \int_{\mathbf{r}_i}^{\mathbf{r}_f} d\mathbf{l} \cdot \mathbf{E}. \quad (4)$$

Hint: Since the electric field is zero inside a perfect conductor, the electric potential inside the conductor must be a constant.

**Solution**