

(Preview of) Final Exam (Fall 2023)

PHYS 500A: MATHEMATICAL METHODS

School of Physics and Applied Physics, Southern Illinois University–Carbondale

Date: 2023 Dec 15

1. (20 points.) Not available in preview mode.
2. (20 points.) Not available in preview mode.
3. (20 points.) The generating function for the Legendre polynomials $P_l(x)$ of degree l is

$$g(t, x) = \frac{1}{\sqrt{1 + t^2 - 2xt}} = \sum_{l=0}^{\infty} t^l P_l(x). \quad (1)$$

- (a) Starting from the generating function and differentiating with respect to t we derived the recurrence relation for Legendre polynomials in Eq. (2),

$$(l + 1) P_{l+1}(x) = (2l + 1)x P_l(x) - l P_{l-1}(x), \quad l = 0, 1, 2, \dots, \quad (2)$$

in terms of

$$P_0(x) = 1 = g(0, x). \quad (3)$$

Differentiating the recurrence relation with respect to x show that

$$(2l + 1) P_l + (2l + 1)x P'_l = l P'_{l-1} + (l + 1) P'_{l+1}, \quad l = 0, 1, 2, \dots, \quad (4)$$

where we suppressed the dependence in x and prime in the superscript of $P'_l(x)$ denotes derivative with respect to the argument x .

- (b) Differentiating the generating function with respect to x show that

$$\frac{\partial g}{\partial x} = \frac{t}{(1 + t^2 - 2xt)^{\frac{3}{2}}} = \sum_{l=0}^{\infty} t^l P'_l(x). \quad (5)$$

Show that the second equality can be rewritten in the form

$$\frac{t}{\sqrt{1 + t^2 - 2xt}} = (1 + t^2 - 2xt) \sum_{l=0}^{\infty} t^l P'_l(x), \quad (6)$$

and implies

$$t \sum_{l=0}^{\infty} t^l P_l(x) = (1 + t^2 - 2xt) \sum_{l=0}^{\infty} t^l P'_l(x). \quad (7)$$

Express this in the form

$$t^0 [P'_0(x)] + t^1 [P'_1(x) - 2xP'_0(x) - P_0(x)] + \sum_{l=2}^{\infty} t^l [P'_l(x) + P'_{l-2}(x) - 2x P'_{l-1}(x) - P_{l-1}(x)] = 0. \quad (8)$$

Then, using the completeness property of Taylor expansion, that is, equating the coefficients of powers of t in the expansion, show that, for t^0 and t^1 ,

$$P'_0(x) = 0, \quad (9a)$$

$$P'_1(x) = P_0(x) = 1, \quad (9b)$$

and matching powers of t^l for $l \geq 2$ derive a recurrence relation for the derivative of Legendre polynomials as

$$2x P'_{l-1} + P_{l-1} = P'_l + P'_{l-2}, \quad l = 2, 3, \dots \quad (10)$$

Here, we shall find it convenient to use the above recurrence relations in the form

$$2x P'_l + P_l = P'_{l+1} + P'_{l-1}, \quad l = 1, 2, 3, \dots, \quad (11)$$

which is obtained by setting $l \rightarrow l + 1$.

- (c) Equations (4) and (11) are linear set of equations for P'_{l-1} and P'_{l+1} in terms of P_l and P'_l . Solve them to find

$$P'_{l+1} = xP'_l + (l+1)P_l, \quad l = 0, 1, 2, \dots, \quad (12a)$$

$$P'_{l-1} = xP'_l - lP_l. \quad l = 1, 2, 3, \dots \quad (12b)$$

- (d) Using $l \rightarrow l - 1$ in Eq. (12a) show that

$$P'_l = xP'_{l-1} + lP_{l-1}. \quad (13)$$

Then, substitute Eq. (12b) to obtain

$$(1-x^2)P'_l = lP_{l-1} - xlP_l. \quad (14)$$

Differentiate the above equation and substitute Eq. (12b) again to derive the differential equation for Legendre polynomials as

$$\left[\frac{\partial}{\partial x} (1-x^2) \frac{\partial}{\partial x} + l(l+1) \right] P_l(x) = 0. \quad (15)$$

Substitute $x = \cos \theta$ to rewrite the differential equation in the form

$$\left[\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \sin \theta \frac{\partial}{\partial \theta} + l(l+1) \right] P_l(\cos \theta) = 0. \quad (16)$$

4. (20 points.) The generating function for the Legendre polynomials $P_l(x)$ of degree l is

$$g(t, x) = \frac{1}{\sqrt{1+t^2-2xt}} = \sum_{l=0}^{\infty} t^l P_l(x). \quad (17)$$

(a) Using binomial expansion show that

$$\frac{1}{\sqrt{1-y}} = \sum_{m=0}^{\infty} y^m \frac{(2m)!}{[m! 2^m]^2} \quad (18)$$

and

$$(2xt - t^2)^m = \sum_{n=0}^{\infty} \frac{m!}{n!(m-n)!} (2xt)^{m-n} t^{2n} (-1)^n. \quad (19)$$

Thus, show that

$$\frac{1}{\sqrt{1+t^2-2xt}} = \sum_{m=0}^{\infty} \sum_{n=0}^m t^{m+n} \frac{(2m)!}{m!n!(m-n)!2^{m+n}} x^{m-n} (-1)^n. \quad (20)$$




Figure 1: Double summation.

(b) In Figure 1 we illustrate how we change the double sum in m and n to variables l and s . This is achieved using the substitutions

$$m + n = l, \quad (21a)$$

$$m - n = 2s, \quad (21b)$$

which corresponds to

$$2m = l + 2s, \quad m = \frac{l}{2} + s, \quad \text{and} \quad n = \frac{l}{2} - s. \quad (22)$$

The counting on the variable s , for given l , follows the pattern,

$$l \text{ even : } 2s = 0, 2, 4, \dots, l, \quad (23a)$$

$$l \text{ odd : } 2s = 1, 3, 5, \dots, l. \quad (23b)$$

Show that in terms of l and s the double summation can be expressed as

$$\frac{1}{\sqrt{1+t^2-2xt}} = \sum_{l=0}^{\infty} \sum_s t^l \frac{(l+2s)!}{\left(\frac{l}{2}+s\right)! \left(\frac{l}{2}-s\right)! (2s)! 2^l} x^{2s} (-1)^{\frac{l}{2}-s}, \quad (24)$$

where the limits on the sum in s are dictated by Eqs. (23) depending on l being even or odd. Thus, read out the polynomial expression for Legendre polynomials of degree l to be

$$P_l(x) = \sum_s \frac{(l+2s)!}{\left(\frac{l}{2}+s\right)! \left(\frac{l}{2}-s\right)! (2s)! 2^l} x^{2s} (-1)^{\frac{l}{2}-s}, \quad (25)$$

where the summation on s depends on whether l is even or odd.

(c) Show that

$$\left(\frac{d}{dx}\right)^l x^{l+2s} = \frac{(l+2s)!}{(2s)!} x^{2s}. \quad (26)$$

Thus, show that

$$P_l(x) = \frac{1}{l! 2^l} \left(\frac{d}{dx}\right)^l \sum_s \frac{l!}{\left(\frac{l}{2}+s\right)! \left(\frac{l}{2}-s\right)!} x^{l+2s} (-1)^{\frac{l}{2}-s}. \quad (27)$$

(d) For even l the summation in s runs from $s = 0$ to $s = l/2$, Thus, writing $l + 2s = 2[l - (\frac{l}{2} - s)]$, show that

$$P_l(x) = \frac{1}{l! 2^l} \left(\frac{d}{dx}\right)^l \sum_{s=0}^{\frac{l}{2}} \frac{l!}{\left(\frac{l}{2}+s\right)! \left(\frac{l}{2}-s\right)!} (x^2)^{l-(\frac{l}{2}-s)} (-1)^{\left(\frac{l}{2}-s\right)}. \quad (28)$$

Then, substituting

$$\frac{l}{2} - s = n, \quad (29)$$

show that

$$P_l(x) = \frac{1}{l! 2^l} \left(\frac{d}{dx}\right)^l \sum_{n=0}^{\frac{l}{2}} \frac{l!}{(l-n)! n!} (x^2)^{l-n} (-1)^n. \quad (30)$$

Note that the summation on n runs from $n = 0$ to $n = l/2$. If we were to extend this sum to $n = l$ verify that the additional terms will have powers in x less than l . Since the terms in the sum are acted upon by l derivatives with respect to x these additional terms will not contribute. Thus, show that

$$P_l(x) = \left(\frac{d}{dx}\right)^l \frac{(x^2 - 1)^l}{l! 2^l}. \quad (31)$$

Similarly, for odd l the summation is s runs as

$$2s = 1, 3, 5, \dots, l, \quad (32)$$

or

$$\frac{2s - 1}{2} = 0, 1, 2, \dots, \frac{l - 1}{2}. \quad (33)$$

Thus, substituting

$$s' = \frac{2s - 1}{2} = s - \frac{1}{2}, \quad (34)$$

show that

$$P_l(x) = \frac{1}{l! 2^l} \left(\frac{d}{dx}\right)^l \sum_{s=0}^{\frac{l-1}{2}} \frac{l!}{\left(\frac{l+1}{2} + s\right)! \left(\frac{l-1}{2} - s\right)!} x^{l+1+2s} (-1)^{\left(\frac{l-1}{2} - s\right)}. \quad (35)$$

Substituting

$$\frac{l - 1}{2} - s = n \quad (36)$$

and writing

$$\frac{l + 1}{2} + s = l - \left(\frac{l - 1}{2} - s\right) \quad (37)$$

show that

$$P_l(x) = \frac{1}{l! 2^l} \left(\frac{d}{dx}\right)^l \sum_{n=0}^{\frac{l-1}{2}} \frac{l!}{(l - n)! n!} (x^2)^{l-n} (-1)^n. \quad (38)$$

Again, like in the case of even l we can extend the sum on n beyond $n = (l - 1)/2$, because they do not survive under the action of l derivatives with respect to x . Thus, again, we have

$$P_l(x) = \left(\frac{d}{dx}\right)^l \frac{(x^2 - 1)^l}{l! 2^l}, \quad (39)$$

which is exactly the form obtained for even l . The expression in Eq.(39) is the Rodrigues formula for generating the Legendre polynomials of degree l .

5. **(20 points.)** Consider the electric potential due to a solid sphere with uniform charge density Q . The angular integral in this evaluation involves the integral

$$\frac{1}{2} \int_{-1}^1 dt \frac{1}{\sqrt{r^2 + r'^2 - 2rr't}}. \quad (40)$$

Evaluate the integral for $r < r'$ and $r' < r$, where r and r' are distances measured from the center of the sphere. (Hint: Substitute $r^2 + r'^2 - 2rr't = y$.)