

Homework No. 07 (Fall 2023)

PHYS 500A: MATHEMATICAL METHODS

School of Physics and Applied Physics, Southern Illinois University–Carbondale

Due date: Monday, 2023 Oct 16, 4.30pm

1. **(20 points.)** Recall that analytic functions satisfy the Cauchy-Riemann equations. That is, the real and imaginary parts of an analytic function

$$f(x + iy) = u(x, y) + iv(x, y) \quad (1)$$

satisfy

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad (2a)$$

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}. \quad (2b)$$

Given $f(z)$ and $g(z)$ are analytic functions in a region, then show that $f(g(z))$ satisfies the Cauchy-Riemann equations there.

Hint: Let $g = u + iv$ and $f = U + iV$. Thus, we can write

$$f(g(z)) = U(u(x, y), v(x, y)) + iV(u(x, y), v(x, y)). \quad (3)$$

2. **(20 points.)** Given an analytic function

$$f(z) = u(x, y) + iv(x, y) \quad (4)$$

and the gradient operator

$$\nabla = \hat{\mathbf{i}} \frac{\partial}{\partial x} + \hat{\mathbf{j}} \frac{\partial}{\partial y}. \quad (5)$$

Show that Cauchy-Riemann equations imply

$$(\nabla u) \cdot (\nabla v) = 0. \quad (6)$$

Thus, interpret that u 's and v 's are orthogonal family of surfaces at every point.

3. **(20 points.)** Analytic functions are significantly constrained, in that they have to satisfy the Cauchy-Riemann conditions. These conditions are necessary (but not sufficient) for a function of a complex variable to be analytic (differentiable). Check if the following functions satisfy the Cauchy-Riemann conditions. If $f(z)$ is analytic for all z , then report

the derivative as a function of z . Otherwise, determine the points, or regions, in the z plane where the function is not analytic.

$$f(z) = z^3, \quad (7a)$$

$$f(z) = |z|^2, \quad (7b)$$

$$f(z) = e^{iz}, \quad (7c)$$

$$f(z) = \ln z, \quad (7d)$$

$$f(z) = e^z + e^{iz}. \quad (7e)$$

4. **(20 points.)** Check if the function

$$f(z) = zz^* \quad (8)$$

satisfies the Cauchy-Riemann conditions.

- (a) Verify that all the points for $f(z)$ lies on the non-negative real line.
- (b) Verify that as you approach the point $z = r \geq 0$ on the non-negative real line, along a circle of fixed radius r from the first quadrant, we have

$$\lim_{\Delta z \rightarrow 0} \frac{\Delta f}{\Delta z} = \lim_{\theta \rightarrow 0} \frac{f(re^{i\theta}) - f(r)}{re^{i\theta} - r} = 0. \quad (9)$$

Then verify that as you approach the point $z = r$ along the real axis we have

$$\lim_{\Delta z \rightarrow 0} \frac{\Delta f}{\Delta z} = \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^2 + y^2 - (x^2 + y^2)}{\Delta x} = 2x. \quad (10)$$

- (c) Thus, conclude that the derivative is not isotropic for any z .

5. **(20 points.)** Check if the function

$$f(z) = \frac{1}{z} \quad (11)$$

satisfies the Cauchy-Riemann conditions.

- (a) Verify that the Cauchy-Riemann conditions for this case are not well defined at $z = 0$, but are fine for $z \neq 0$.
- (b) Verify that

$$\frac{df}{dz} = -\frac{1}{z^2}, \quad z \neq 0. \quad (12)$$

- (c) Determine the limiting value of the derivative as you approach $z = 0$ along the positive real line, and, then, when you approach along the negative real line. Repeat the analysis along the imaginary line. Repeat the analysis along the line $x = y$. Are these limits identical?
- (d) If these limits are not identical conclude that the derivative is not isotropic at $z = 0$. Then, the function is not analytic at $z = 0$.