

Homework No. 03 (Fall 2024)

PHYS 500A: MATHEMATICAL METHODS

School of Physics and Applied Physics, Southern Illinois University–Carbondale

Due date: Friday, 2024 Sep 13, 4.30pm

0. Problems 1 and 4 are for practice. Problem 2 and 3 are for submission.

1. (**Example.**) Let \mathbf{r} represent the position vector, x^i the components of the position vector in rectangular coordinates, and u^i the components of the position vector in cylindrical polar coordinates. In particular, we have

$$x^1 = x = \rho \cos \phi, \quad u^1 = \rho = \sqrt{x^2 + y^2}, \quad (1a)$$

$$x^2 = y = \rho \sin \phi, \quad u^2 = \phi = \tan^{-1} \frac{y}{x}, \quad (1b)$$

$$x^3 = z = z, \quad u^3 = z = z. \quad (1c)$$

Let us define the unit vectors

$$\hat{\rho} = \cos \phi \hat{\mathbf{i}} + \sin \phi \hat{\mathbf{j}} + 0 \hat{\mathbf{k}}, \quad (2a)$$

$$\hat{\phi} = -\sin \phi \hat{\mathbf{i}} + \cos \phi \hat{\mathbf{j}} + 0 \hat{\mathbf{k}}, \quad (2b)$$

$$\hat{\mathbf{z}} = 0 \hat{\mathbf{i}} + 0 \hat{\mathbf{j}} + \hat{\mathbf{k}}, \quad (2c)$$

where $\hat{\mathbf{i}}$, $\hat{\mathbf{j}}$, and $\hat{\mathbf{k}}$ are basis vectors in rectangular coordinate system. We will also use the notation $\hat{\mathbf{i}} = \hat{\mathbf{x}}^1 = \hat{\mathbf{x}}_1$, $\hat{\mathbf{j}} = \hat{\mathbf{x}}^2 = \hat{\mathbf{x}}_2$, $\hat{\mathbf{k}} = \hat{\mathbf{x}}^3 = \hat{\mathbf{x}}_3$, to represent these vectors.

(a) Basis vectors:

$$\mathbf{e}_i = \frac{\partial \mathbf{r}}{\partial u^i}. \quad (3)$$

Show that

$$\mathbf{e}_1 = \hat{\rho}, \quad \mathbf{e}_2 = \rho \hat{\phi}, \quad \mathbf{e}_3 = \hat{\mathbf{z}}. \quad (4)$$

(b) Reciprocal basis vectors:

$$\mathbf{e}^i = \nabla u^i. \quad (5)$$

Show that

$$\mathbf{e}^1 = \hat{\rho}, \quad \mathbf{e}^2 = \frac{\hat{\phi}}{\rho}, \quad \mathbf{e}^3 = \hat{\mathbf{z}}. \quad (6)$$

(c) Orthonormality: Show that

$$\mathbf{e}^i \cdot \mathbf{e}_j = \delta_j^i. \quad (7)$$

(d) Metric tensor: A line element is defined as

$$d\mathbf{r} = dx^i \hat{\mathbf{x}}_i = du^i \mathbf{e}_i. \quad (8)$$

Show that

$$d\mathbf{r} \cdot d\mathbf{r} = du^i du^j g_{ij}, \quad (9)$$

where the metric tensor g_{ij} is defined as

$$g_{ij} = \mathbf{e}_i \cdot \mathbf{e}_j. \quad (10)$$

Evaluate all the components of g_{ij} .

(e) Completeness relation: Starting from

$$\nabla \mathbf{r} = \mathbf{1} \quad (11)$$

derive the completeness relation

$$\mathbf{e}^i \mathbf{e}_i = \mathbf{1}. \quad (12)$$

Express the completeness relation in cylindrical polar coordinates in terms of $\hat{\boldsymbol{\rho}}$, $\hat{\boldsymbol{\phi}}$, and $\hat{\mathbf{z}}$.

(f) Transformation matrix: The components of a vector \mathbf{A} are defined using the relations

$$\mathbf{A} = \mathbf{A} \cdot \mathbf{1} = A^i \hat{\mathbf{x}}_i = \bar{A}^i \mathbf{e}_i, \quad (13a)$$

$$\mathbf{A} = \mathbf{1} \cdot \mathbf{A} = \hat{\mathbf{x}}^i A_i = \mathbf{e}^i \bar{A}_i. \quad (13b)$$

Then, derive the transformations

$$\bar{A}^j = A^i T_i^j, \quad T_i^j = \hat{\mathbf{x}}_i \cdot \mathbf{e}^j, \quad (14a)$$

$$\bar{A}_j = S_j^i A_i, \quad S_j^i = \mathbf{e}_j \cdot \hat{\mathbf{x}}^i, \quad (14b)$$

and show that $T_i^j S_j^k = \delta_i^k$. Find S and T for cylindrical polar coordinates.

2. (**Example.**) Let \mathbf{r} represent the position vector, x^i the components of the position vector in rectangular coordinates, and u^i the components of the position vector in spherical polar coordinates. In particular, we have

$$x^1 = x = r \sin \theta \cos \phi, \quad u^1 = r = \sqrt{x^2 + y^2 + z^2}, \quad (15a)$$

$$x^2 = y = r \sin \theta \sin \phi, \quad u^2 = \theta = \tan^{-1} \frac{\sqrt{x^2 + y^2}}{z}, \quad (15b)$$

$$x^3 = z = r \cos \theta, \quad u^3 = \phi = \tan^{-1} \frac{y}{x}. \quad (15c)$$

Let us define the unit vectors

$$\hat{\mathbf{r}} = \sin \theta \cos \phi \hat{\mathbf{i}} + \sin \theta \sin \phi \hat{\mathbf{j}} + \cos \theta \hat{\mathbf{k}}, \quad (16a)$$

$$\hat{\boldsymbol{\theta}} = \cos \theta \cos \phi \hat{\mathbf{i}} + \cos \theta \sin \phi \hat{\mathbf{j}} - \sin \theta \hat{\mathbf{k}}, \quad (16b)$$

$$\hat{\boldsymbol{\phi}} = -\sin \phi \hat{\mathbf{i}} + \cos \phi \hat{\mathbf{j}}, \quad (16c)$$

where $\hat{\mathbf{i}}$, $\hat{\mathbf{j}}$, and $\hat{\mathbf{k}}$ are basis vectors in rectangular coordinate system. We will also use the notation $\hat{\mathbf{i}} = \hat{\mathbf{x}}^1 = \hat{\mathbf{x}}_1$, $\hat{\mathbf{j}} = \hat{\mathbf{x}}^2 = \hat{\mathbf{x}}_2$, $\hat{\mathbf{k}} = \hat{\mathbf{x}}^3 = \hat{\mathbf{x}}_3$, to represent these vectors.

(a) Basis vectors:

$$\mathbf{e}_i = \frac{\partial \mathbf{r}}{\partial u^i}. \quad (17)$$

Show that

$$\mathbf{e}_1 = \hat{\mathbf{r}}, \quad \mathbf{e}_2 = r\hat{\boldsymbol{\theta}}, \quad \mathbf{e}_3 = r \sin \theta \hat{\boldsymbol{\phi}}. \quad (18)$$

(b) Reciprocal basis vectors:

$$\mathbf{e}^i = \nabla u^i. \quad (19)$$

Show that

$$\mathbf{e}^1 = \hat{\mathbf{r}}, \quad \mathbf{e}^2 = \frac{\hat{\boldsymbol{\theta}}}{r}, \quad \mathbf{e}^3 = \frac{\hat{\boldsymbol{\phi}}}{r \sin \theta}. \quad (20)$$

(c) Orthonormality: Show that

$$\mathbf{e}^i \cdot \mathbf{e}_j = \delta_j^i. \quad (21)$$

(d) Metric tensor: A line element is defined as

$$d\mathbf{r} = dx^i \hat{\mathbf{x}}_i = du^i \mathbf{e}_i. \quad (22)$$

Show that

$$d\mathbf{r} \cdot d\mathbf{r} = du^i du^j g_{ij}, \quad (23)$$

where the metric tensor g_{ij} is defined as

$$g_{ij} = \mathbf{e}_i \cdot \mathbf{e}_j. \quad (24)$$

Evaluate all the components of g_{ij} .

(e) Completeness relation: Starting from

$$\nabla \mathbf{r} = \mathbf{1} \quad (25)$$

derive the completeness relation

$$\mathbf{e}^i \mathbf{e}_i = \mathbf{1}. \quad (26)$$

Express the completeness relation in spherical polar coordinates in terms of $\hat{\mathbf{r}}$, $\hat{\boldsymbol{\theta}}$, and $\hat{\boldsymbol{\phi}}$.

(f) Transformation matrix: The components of a vector \mathbf{A} are defined using the relations

$$\mathbf{A} = \mathbf{A} \cdot \mathbf{1} = A^i \hat{\mathbf{x}}_i = \bar{A}^i \mathbf{e}_i, \quad (27a)$$

$$\mathbf{A} = \mathbf{1} \cdot \mathbf{A} = \hat{\mathbf{x}}^i A_i = \mathbf{e}^i \bar{A}_i. \quad (27b)$$

Then, derive the transformations

$$\bar{A}^j = A^i T_i^j, \quad T_i^j = \hat{\mathbf{x}}_i \cdot \mathbf{e}^j, \quad (28a)$$

$$\bar{A}_j = S_j^i A_i, \quad S_j^i = \mathbf{e}_j \cdot \hat{\mathbf{x}}^i, \quad (28b)$$

and show that $T_i^j S_j^k = \delta_i^k$. Find S and T for spherical polar coordinates.

3. (20 points.) Let \mathbf{r} represent a position vector in three dimensional space. Let x^i be the components of the position vector in rectangular coordinates, which can be interpreted as surfaces of constant x^i . Let us coordinatize the space using the planes, labeled using β ,

$$y = mx + \beta \quad (29)$$

where m is fixed, instead of planes with constant y . The other two sets of planes of constant x and constant z are the same. See Fig. 1. Let u^i be the components of the position vector in this new coordinatization of space. In particular, we have




Figure 1: Basis vectors \mathbf{e}_i and reciprocal basis vectors \mathbf{e}^i .

$$x^1 = x = \alpha, \quad u^1 = \alpha = x, \quad (30a)$$

$$x^2 = y = mx + \beta, \quad u^2 = \beta = y - mx, \quad (30b)$$

$$x^3 = z = \gamma, \quad u^3 = \gamma = z. \quad (30c)$$

The basis vectors $\hat{\mathbf{i}}$, $\hat{\mathbf{j}}$, and $\hat{\mathbf{k}}$ in rectangular coordinate system will be represented as $\hat{\mathbf{i}} = \hat{\mathbf{x}}^1 = \hat{\mathbf{x}}_1$, $\hat{\mathbf{j}} = \hat{\mathbf{x}}^2 = \hat{\mathbf{x}}_2$, $\hat{\mathbf{k}} = \hat{\mathbf{x}}^3 = \hat{\mathbf{x}}_3$, if necessary.

- (a) Basis vectors:

$$\mathbf{e}_i = \frac{\partial \mathbf{r}}{\partial u^i}. \quad (31)$$

Show that

$$\mathbf{e}_1 = \hat{\mathbf{i}} + m\hat{\mathbf{j}}, \quad \mathbf{e}_2 = \hat{\mathbf{j}}, \quad \mathbf{e}_3 = \hat{\mathbf{k}}. \quad (32)$$

- (b) Reciprocal basis vectors:

$$\mathbf{e}^i = \nabla u^i. \quad (33)$$

Show that

$$\mathbf{e}^1 = \hat{\mathbf{i}}, \quad \mathbf{e}^2 = -m\hat{\mathbf{i}} + \hat{\mathbf{j}}, \quad \mathbf{e}^3 = \hat{\mathbf{k}}. \quad (34)$$

Verify the relations

$$\mathbf{e}^1 = \frac{\mathbf{e}_2 \times \mathbf{e}_3}{(\mathbf{e}_2 \times \mathbf{e}_3) \cdot \mathbf{e}_1}, \quad \mathbf{e}^2 = \frac{\mathbf{e}_3 \times \mathbf{e}_1}{(\mathbf{e}_3 \times \mathbf{e}_1) \cdot \mathbf{e}_2}, \quad \mathbf{e}^3 = \frac{\mathbf{e}_1 \times \mathbf{e}_2}{(\mathbf{e}_1 \times \mathbf{e}_2) \cdot \mathbf{e}_3}. \quad (35)$$

(c) Orthonormality: Show that

$$\mathbf{e}^i \cdot \mathbf{e}_j = \delta_j^i. \quad (36)$$

That is,

$$\mathbf{e}^1 \cdot \mathbf{e}_1 = 1, \quad \mathbf{e}^1 \cdot \mathbf{e}_2 = 0, \quad \mathbf{e}^1 \cdot \mathbf{e}_3 = 0, \quad (37a)$$

$$\mathbf{e}^2 \cdot \mathbf{e}_1 = 0, \quad \mathbf{e}^2 \cdot \mathbf{e}_2 = 1, \quad \mathbf{e}^2 \cdot \mathbf{e}_3 = 0, \quad (37b)$$

$$\mathbf{e}^3 \cdot \mathbf{e}_1 = 0, \quad \mathbf{e}^3 \cdot \mathbf{e}_2 = 0, \quad \mathbf{e}^3 \cdot \mathbf{e}_3 = 1. \quad (37c)$$

(d) Metric tensor: The metric tensor g_{ij} is defined as

$$g_{ij} = \mathbf{e}_i \cdot \mathbf{e}_j. \quad (38)$$

Evaluate all the components of g_{ij} . That is,

$$g_{11} = \mathbf{e}_1 \cdot \mathbf{e}_1 = 1 + m^2, \quad g_{12} = \mathbf{e}_1 \cdot \mathbf{e}_2 = m, \quad g_{13} = \mathbf{e}_1 \cdot \mathbf{e}_3 = 0, \quad (39a)$$

$$g_{21} = \mathbf{e}_2 \cdot \mathbf{e}_1 = m, \quad g_{22} = \mathbf{e}_2 \cdot \mathbf{e}_2 = 1, \quad g_{23} = \mathbf{e}_2 \cdot \mathbf{e}_3 = 0, \quad (39b)$$

$$g_{31} = \mathbf{e}_3 \cdot \mathbf{e}_1 = 0, \quad g_{32} = \mathbf{e}_3 \cdot \mathbf{e}_2 = 0, \quad g_{33} = \mathbf{e}_3 \cdot \mathbf{e}_3 = 1. \quad (39c)$$

Similarly evaluate the components of

$$g^{ij} = \mathbf{e}^i \cdot \mathbf{e}^j. \quad (40)$$

That is,

$$g^{11} = \mathbf{e}^1 \cdot \mathbf{e}^1 = 1, \quad g^{12} = \mathbf{e}^1 \cdot \mathbf{e}^2 = -m, \quad g^{13} = \mathbf{e}^1 \cdot \mathbf{e}^3 = 0, \quad (41a)$$

$$g^{21} = \mathbf{e}^2 \cdot \mathbf{e}^1 = -m, \quad g^{22} = \mathbf{e}^2 \cdot \mathbf{e}^2 = 1 + m^2, \quad g^{23} = \mathbf{e}^2 \cdot \mathbf{e}^3 = 0, \quad (41b)$$

$$g^{31} = \mathbf{e}^3 \cdot \mathbf{e}^1 = 0, \quad g^{32} = \mathbf{e}^3 \cdot \mathbf{e}^2 = 0, \quad g^{33} = \mathbf{e}^3 \cdot \mathbf{e}^3 = 1. \quad (41c)$$

Verify that $g^{ij}g_{jk} = \delta_k^i$.

(e) Completeness relation: Verify the completeness relation

$$\mathbf{e}^i \mathbf{e}_i = \mathbf{1} \quad (42)$$

by evaluating

$$\mathbf{e}^1 \mathbf{e}_1 + \mathbf{e}^2 \mathbf{e}_2 + \mathbf{e}^3 \mathbf{e}_3. \quad (43)$$

(f) Given a vector

$$\mathbf{A} = a \hat{\mathbf{i}} + b \hat{\mathbf{j}} + c \hat{\mathbf{k}} \quad (44)$$

in rectangular coordinates, find the components of the vector \mathbf{A} in the basis of \mathbf{e}_i . That is, find the components A^i in

$$\mathbf{A} = A^1 \mathbf{e}_1 + A^2 \mathbf{e}_2 + A^3 \mathbf{e}_3. \quad (45)$$

4. (20 points.) The tangent and normal vectors for the cylindrical coordinate system are

$$\mathbf{e}_1 = \mathbf{e}_\rho = \hat{\boldsymbol{\rho}}, \quad \mathbf{e}^1 = \mathbf{e}^\rho = \hat{\boldsymbol{\rho}}, \quad (46a)$$

$$\mathbf{e}_2 = \mathbf{e}_\phi = \rho \hat{\boldsymbol{\phi}}, \quad \mathbf{e}^2 = \mathbf{e}^\phi = \frac{\hat{\boldsymbol{\phi}}}{\rho}, \quad (46b)$$

$$\mathbf{e}_3 = \mathbf{e}_z = \hat{\mathbf{z}}, \quad \mathbf{e}^3 = \mathbf{e}^z = \hat{\mathbf{z}}. \quad (46c)$$

The connection is defined as

$$(\nabla \mathbf{e}_i). \quad (47)$$

Berry connection $\mathbf{A}_i{}^k$ captures the projections of the connection to the right,

$$\mathbf{A}_i{}^k = (\nabla \mathbf{e}_i) \cdot \mathbf{e}^k. \quad (48)$$

Compute the Berry connection $\mathbf{A}_i{}^k$ for the cylindrical coordinate system to be

$$\mathbf{A}_i{}^k = \begin{pmatrix} 0 & \frac{\hat{\boldsymbol{\phi}}}{\rho^2} & 0 \\ -\hat{\boldsymbol{\phi}} & \frac{\boldsymbol{\rho}}{\rho^2} & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (49)$$

The Christoffel symbols Γ_{ij}^k captures all the projections of the connection,

$$\mathbf{e}_j \cdot (\nabla \mathbf{e}_i) \cdot \mathbf{e}^k = \Gamma_{ij}^k. \quad (50)$$

Compute the Christoffel symbols for the cylindrical coordinate system. Show that the non-zero Christoffel symbols in cylindrical coordinates are

$$\Gamma_{22}^1 = \Gamma_{\phi\phi}^\rho = -\rho, \quad (51)$$

$$\Gamma_{12}^2 = \Gamma_{\rho\phi}^\phi = \frac{1}{\rho}, \quad (52)$$

$$\Gamma_{21}^2 = \Gamma_{\phi\rho}^\phi = \frac{1}{\rho}. \quad (53)$$