

Homework No. 04 (2026 Spring)

PHYS 510: CLASSICAL MECHANICS

School of Physics and Applied Physics, Southern Illinois University–Carbondale

Due date: Thursday, 2026 Feb 12, 4.30pm

1. (**20 points.**) The motion of a particle of mass m near the Earth's surface is described by

$$\frac{d}{dt}(mv) = -mg, \quad (1)$$

where $v = dz/dt$ is the velocity in the upward z direction.

- (a) Find the Lagrangian for this system that implies the equation of motion of Eq. (1) using the principle of stationary action.
 - (b) Determine the canonical momentum for this system
 - (c) Determine the Hamilton $H(p, z)$ for this system.
 - (d) Determine the Hamilton equations of motion.
2. (**20 points.**) The motion of a particle of mass m undergoing simple harmonic motion is described by

$$\frac{d}{dt}(mv) = -kx, \quad (2)$$

where $v = dx/dt$ is the velocity in the x direction.

- (a) Find the Lagrangian for this system that implies the equation of motion of Eq. (2) using the principle of stationary action.
 - (b) Determine the canonical momentum for this system
 - (c) Determine the Hamiltonian $H(p, x)$ for this system.
 - (d) Determine the Hamilton equations of motion.
3. (**20 points.**) (Refer Goldstein, 2nd edition, Chapter 1 Problem 8.) As a consequence of the Hamilton's stationary action principle, the equations of motion for a system can be expressed as Euler-Lagrange equations,

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{x}} - \frac{\partial L}{\partial x} = 0, \quad (3)$$

in terms of a Lagrangian $L(x, \dot{x}, t)$. Show that the Lagrangian for a system is not unique. In particular, show that if $L(x, \dot{x}, t)$ satisfies the Euler-Lagrange equation then

$$L'(x, \dot{x}, t) = L(x, \dot{x}, t) + \frac{dF(x, t)}{dt}, \quad (4)$$

where $F(x, t)$ is any arbitrary differentiable function, also satisfies the Euler-Lagrange equation.

4. (**20 points.**) Consider a (time independent) Hamiltonian

$$H = H(x, p), \quad (5)$$

which satisfies the Hamilton equations of motion

$$\frac{dx}{dt} = \frac{\partial H}{\partial p}, \quad (6a)$$

$$\frac{dp}{dt} = -\frac{\partial H}{\partial x}. \quad (6b)$$

- (a) Recall that the Lagrangian, which will temporarily be called the x -Lagrangian here, is defined by the construction

$$L_x(x, \dot{x}) = p\dot{x} - H(x, p). \quad (7)$$

Starting from Eq. (7) derive

$$\frac{\partial L_x}{\partial x} = -\frac{\partial H}{\partial x}, \quad (8a)$$

$$\frac{\partial L_x}{\partial \dot{x}} = p, \quad (8b)$$

$$\frac{\partial L_x}{\partial p} = \dot{x} - \frac{\partial H}{\partial p}. \quad (8c)$$

Using the Hamilton equations of motion, Eqs. (6), in Eqs. (8) we have the equations governing the x -Lagrangian to be

$$\frac{\partial L_x}{\partial p} = 0, \quad (9a)$$

$$\frac{\partial L_x}{\partial \dot{x}} = p, \quad (9b)$$

$$\frac{d}{dt} \frac{\partial L_x}{\partial \dot{x}} = \frac{\partial L_x}{\partial x}. \quad (9c)$$

- (b) Now, define the p -Lagrangian using the construction

$$L_p(p, \dot{p}) = -x\dot{p} - H(x, p). \quad (10)$$

The opposite sign in the construction of the p -Lagrangian is motivated by the action principle, which does not care for a total derivative, refer Schwinger. Starting from Eq. (10) derive

$$\frac{\partial L_p}{\partial p} = -\frac{\partial H}{\partial p}, \quad (11a)$$

$$\frac{\partial L_p}{\partial \dot{p}} = -x, \quad (11b)$$

$$\frac{\partial L_p}{\partial x} = -\dot{p} - \frac{\partial H}{\partial x}. \quad (11c)$$

Using the Hamilton equations of motion, Eqs. (6), in Eqs. (11) we have the equations governing the p -Lagrangian to be

$$\frac{\partial L_p}{\partial x} = 0, \quad (12a)$$

$$\frac{\partial L_p}{\partial \dot{p}} = -x, \quad (12b)$$

$$\frac{d}{dt} \frac{\partial L_p}{\partial \dot{p}} = \frac{\partial L_p}{\partial p}. \quad (12c)$$

- (c) Illustrate the above using a specific Hamiltonian, for example that of a harmonic oscillator, as a guide.